

# Pediatric Otolaryngology

## Nine Things Clinicians and Patients Should Question

by  
Canadian Society of Otolaryngology–Head & Neck Surgery  
Pediatric Otolaryngology Subspecialty Interest Group

Last updated: December 2020



### 1 **Don't routinely order a plain film x-ray in the evaluation of pediatric nasal fractures.**

Nasal fractures are one of the most common facial fractures in the pediatric population. The decision to perform a closed reduction procedure in the operating room is based on factors such as breathing difficulty and external deformity, which are not assessed effectively by x-ray. Plain film x-rays are unable to accurately evaluate nasal fractures given its low sensitivity and specificity, at 72% and 73% respectively. Physical examination is often sufficient to make a diagnosis for children with displaced nasal fractures. Overall, x-rays do not add value to the diagnosis or treatment plan for children with nasal fractures and should not be ordered to avoid their associated costs and radiation exposure.

### 2 **Don't order imaging to distinguish acute bacterial sinusitis from an upper respiratory infection.**

Acute bacterial sinusitis (ABS) is a diagnosis that is made based on clinical criteria and has a low prevalence amongst children presenting with respiratory symptoms. Although a normal radiograph, CT, or MRI can help to rule out ABS, an abnormal result does not confirm the diagnosis. Given that many children will have abnormal imaging due to a viral upper respiratory infection during certain times of the year, combined with the potential for exposure to radiation, routine imaging is not recommended. Instances in which imaging would be warranted include if the child is immunocompromised, or if orbital, central nervous system, or other suppurative complications are present.

The American Academy of Pediatrics recommends diagnosing pediatric ABS when (1) cough, nasal discharge or both are persistent for >10 days without improvement; (2) there is worsening or new onset of cough, nasal discharge, or fever; or (3) there is a severe onset, with a fever greater  $\geq 39^{\circ}\text{C}$ , concurrently with purulent nasal discharge for at least 3 consecutive days.

### 3 **Don't place tympanostomy tubes in most children for a single episode of uncomplicated otitis media with effusion of less than 3 months' duration.**

Although tympanostomy tube insertion can be associated with short-term quality of life improvements, the natural history of otitis media with effusion (OME) is sufficiently favorable and most OME in children will spontaneously resolve within 3 months. Cases of OME which last longer than 3 months are typically chronic in nature, and less likely to resolve without intervention. Limited data exists regarding the efficacy of tympanostomy tube insertion in children with OME for less than 3 months. By delaying the consideration for tympanostomy tube insertion, potentially unnecessary procedures are avoided, along with the associated risks, tube related side effects, and costs. Children excluded from this recommendation include those who have risk factors for developmental difficulties such as trisomy 21, Autism-spectrum disorder, blindness, and permanent hearing loss independent of OME.

### 4 **Don't routinely prescribe intranasal/systemic steroids, antihistamines or decongestants for children with uncomplicated otitis media with effusion.**

In most cases, medical treatment using antihistamines, decongestants, systemic antibiotics and steroids have shown little to no effect on the long-term outcomes of uncomplicated otitis media with effusion (OME) in children. Because of this, and the costs and potential side effects, it is not recommended to prescribe these medical treatments for children with uncomplicated OME. The exception to this would be for children with coexisting conditions in which these medications are indicated for primary management.

### 5 **Don't prescribe oral antibiotics for children with uncomplicated tympanostomy tube otorrhea or uncomplicated acute otitis externa.**

The use of oral antibiotics where they are not necessary can promote antibiotic resistance and increase the risk of opportunistic infections. Topical antibiotics achieve higher concentrations in the ear canal, demonstrate improved patient satisfaction, are associated with fewer adverse events, and are shown to have equal efficacy for treatment of acute tympanostomy tube otorrhea (TTO) and acute otitis externa (AOE) when compared to oral antibiotics. For these reasons, topical antibiotics rather than oral antibiotics should be prescribed as first line treatment for acute uncomplicated TTO and uncomplicated AOE.

**6 Don't prescribe codeine for post-tonsillectomy/adenoidectomy pain relief in children.**

Codeine has been associated with a high rate of adverse drug reactions in children. This includes life-threatening respiratory depression. Appropriate dosing of codeine is challenging due to the genetic heterogeneity amongst patients for the CYP2D6 enzyme, which is responsible for codeine metabolism. Genetic screening of CYP2D6 is not routinely performed and can not reliably identify variations in codeine metabolism rates amongst patients. As such, children who are ultra-fast metabolizers of codeine are placed at increased risk of severe adverse drug reactions. Alternative analgesia should be used post-tonsillectomy/adenoidectomy.

**7 Don't administer perioperative antibiotics for elective tonsillectomy in children.**

Administration of perioperative antibiotics for children undergoing tonsillectomy shows no significant benefits in regard to common post-tonsillectomy morbidities. Overuse of systemic antibiotics increases bacterial resistance and the risk of adverse drug events unnecessarily. These concerns outweigh the reduction in postoperative fever which is the only potential benefit of perioperative antibiotic administration for elective tonsillectomy. Therefore, perioperative antibiotics are not indicated for children undergoing elective tonsillectomy, unless specific indications are present (e.g., cardiac conditions or those with a peritonsillar abscess or acute infection).

**8 Don't perform tonsillectomy for children with uncomplicated recurrent throat infections if there have been fewer than 7 episodes in the past year, 5 episodes in each of the past 2 years, or 3 episodes in each of the last 3 years.**

For children who have a lower number of recurrent throat infections, tonsillectomy has significantly less benefits when compared to those with more frequent infections, and many children with recurrent throat infections naturally improve without intervention. Therefore, where safely possible, avoidance of tonsillectomy for children with lower numbers of acute infections is recommended. This avoids unnecessary tonsillectomy and the costs and complications associated with the procedure (i.e., bleeding, pain, infection). If tonsillectomy is not indicated, children should be closely monitored and reconsidered for tonsillectomy if the infection frequency increases, as they would be less likely to naturally improve, and more likely to benefit from tonsillectomy. Families should be counselled on the limited benefits and potential harms of performing tonsillectomy for children and adolescents with low rates of recurrent throat infections. Shared decision making is of importance when considering tonsillectomy as individual patient and family factors can impact the decision.

**9 Don't perform endoscopic sinus surgery for uncomplicated pediatric chronic rhinosinusitis prior to failure of maximal medical therapy and adenoidectomy.**

While endoscopic sinus surgery (ESS) has been found to be an effective therapy in children with chronic rhinosinusitis, comparable outcomes can be achieved with medical therapy and adenoidectomy. A stepwise approach of medical therapy, progressing to adenoidectomy, then to ESS allows children to be treated with a less invasive and more cost-effective interventions as initial therapy, while saving ESS for those who are refractory to primary interventions. Maximal medical therapy should be exhausted prior to surgical intervention for uncomplicated patients. In cases with complications such as orbital or skull base involvement, ESS can be employed more readily.

## How the list was created

This list was created by the Pediatric Otolaryngology Subspecialty Interest Group of the Canadian Society of Otolaryngology–Head & Neck Surgery. A list of 25 recommendations regarding unnecessary tests and interventions along with evidence supporting them were compiled. These unnecessary tests and interventions are often invasive and incur risk to patients and unwarranted costs to our public health care system. The members of the Subspecialty Interest Group were asked to provide feedback on the recommendations and to rate them regarding five factors: potential to affect clinical practice; cost-benefit ratio; evidence supporting recommendation; pervasiveness of test/intervention; and potential to cause harm. The final list was then selected and edited based on the group members ratings and feedback.

## Sources

- 1 Desrosiers AE 3rd, Thaller SR. Pediatric nasal fractures: evaluation and management. *J Craniofac Surg*. 2011;22(4):1327-9. [PMID: 21772190](#).  
Mohammadi A, Ghasemi-Rad M. Nasal bone fracture--ultrasonography or computed tomography? *Med Ultrason*. 2011;13(4):292-5. [PMID: 22132401](#).  
Nigam A, Goni A, Benjamin A, Dasgupta AR. The value of radiographs in the management of the fractured nose. *Arch Emerg Med*. 1993;10(4):293-7. [PMID: 8110318](#).
- 2 Aitken M, Taylor JA. Prevalence of clinical sinusitis in young children followed up by primary care pediatricians. *Arch Pediatr Adolesc Med*. 1998;152(3):244-8. [PMID: 9529461](#).  
Gwaltney JM Jr, Phillips CD, Miller RD, Riker DK. Computed tomographic study of the common cold. *N Engl J Med*. 1994;330(1):25-30. [PMID: 8259141](#).  
Kristo A, Uhari M, Luotonen J, Koivunen P, Ilkko E, Tapiainen T, et al. Paranasal sinus findings in children during respiratory infection evaluated with magnetic resonance imaging. *Pediatrics*. 2003;111(5 Pt 1):e586-9. [PMID: 12728114](#).  
Wald ER, Applegate KE, Bordley C, Darrow DH, Glode MP, Marcy SM, et al. Clinical practice guideline for the diagnosis and management of acute bacterial sinusitis in children aged 1 to 18 years. *Pediatrics*. 2013;132(1):e262-80. [PMID: 23796742](#).  
Wald ER, Nash D, Eickhoff J. Effectiveness of amoxicillin/clavulanate potassium in the treatment of acute bacterial sinusitis in children. *Pediatrics*. 2009;124(1):9-15. [PMID: 19564277](#).
- 3 Hellstrom S, Groth A, Jorgensen F. Ventilation tube treatment: a systematic review of the literature. *Otolaryngol Head Neck Surg*. 2011;145(3):383-95. [PMID: 21632976](#).  
Lous J, Burton MJ, Felding JU, Ovesen T, Rovers MM, Williamson I. Grommets (ventilation tubes) for hearing loss associated with otitis media with effusion in children. *Cochrane Database Syst Rev*. 2005;(1):CD001801. [PMID: 15674886](#).  
Rosenfeld RM, Kay D. Natural history of untreated otitis media. *Laryngoscope*. 2003;113(10):1645-57. [PMID: 14520089](#).  
Rosenfeld RM, Schwartz SR, Pynnonen MA, Tunkel DE, Hussey HM, Fichera JS, et al. Clinical practice guideline: tympanostomy tubes in children. *Otolaryngol Head Neck Surg*. 2013;149(1):S1-35. [PMID: 23818543](#).
- 4 Griffin, G, Flynn, CA. Antihistamines and/or decongestants for otitis media with effusion (OME) in children. *Cochrane Database Syst Rev*. 2011;9:CD003423. [PMID: 21901683](#).  
Rosenfeld RM, Shin JJ, Schwartz SR, Coggins R, Gagnon L, Hackell JM, et al. Clinical practice guideline: otitis media with effusion (update). *Otolaryngol Head Neck Surg*. 2016;154(1 Suppl):S1-S41. [PMID: 26832942](#).  
Simpson SA, Lewis R, van der Voort J, Butler CC. Oral or topical nasal steroids for hearing loss associated with otitis media with effusion in children. *Cochrane Database Syst Rev*. 2011;(5):CD001935. [PMID: 21563132](#).  
Venekamp RP, Burton MJ, van Dongen TM, van der Heijden GJ, van Zon A, Schilder AG. Antibiotics for otitis media with effusion in children. *Cochrane Database Syst Rev*. 2016;(6):CD009163. [PMID: 27290722](#).
- 5 Goldblatt EL, Dohar J, Nozza RJ, Nielsen RW, Goldberg T, Sidman JD, et al. Topical ofloxacin versus systemic amoxicillin/clavulanate in purulent otorrhea in children with tympanostomy tubes. *Int J Pediatr Otorhinolaryngol*. 1998;46(1-2):91-101. [PMID: 10190709](#). <https://pubmed.ncbi.nlm.nih.gov/10190709/>  
Kaushik V, Malik T, Saeed SR. Interventions for acute otitis externa. *Cochrane Database of Syst Rev*. 2010;6(2):444–560. [PMID: 20091565](#).  
Rosenfeld RM, Schwartz SR, Pynnonen MA, Tunkel DE, Hussey HM, Fichera JS, et al. Clinical practice guideline: tympanostomy tubes in children. *Otolaryngol Head Neck Surg*. 2013;149(1 Suppl):S1-35. [PMID: 23818543](#).  
Rosenfeld RM, Schwartz SR, Cannon CR, Roland PS, Simon GR, Kumar KA, et al. Clinical practice guideline: acute otitis externa. *Otolaryngol Head Neck Surg*. 2014;150(1):S1-S24. [PMID: 24491310](#).  
Rosenfeld RM, Singer M, Wasserman JM, Stinnett SS. Systematic review of topical antimicrobial therapy for acute otitis externa. *Otolaryngol Head Neck Surg*. 2006;134:S24-48. [PMID: 16638474](#).
- 6 Crews KR, Gaedigk A, Dunnenberger HM, et al. Clinical pharmacogenetics implementation consortium guidelines for cytochrome P450 2D6 genotype and codeine therapy: 2014 update. *Clin Pharmacol Ther*. 2014;95(4):376-82. [PMID: 24458010](#).  
Kelly LE, Rieder M, van den Anker J, Malkin B, Ross C, Neely MN, et al. More codeine fatalities after tonsillectomy in North American children. *Pediatrics*. 2012;129(5):e1343-7. [PMID: 22492761](#).  
Mitchell RB, Archer SM, Ishman SL, Rosenfeld RM, Coles S, Finestone SA, et al. Clinical practice guideline: tonsillectomy in children (update). *Otolaryngol Head Neck Surg*. 2019;160(1 Suppl):S1–42. [PMID: 30921525](#).  
Prows CA, Zhang X, Huth MM, Zhang K, Saldaña SN, Daraiseh NM, et al. Codeine-related adverse drug reactions in children following tonsillectomy: a prospective study. *Laryngoscope*. 2014;124(5):1242–50. [PMID: 24122716](#).  
Tobias JD, Green TP, Coté CJ. Codeine: time to say no. *Pediatrics*. 2016;138(4):e20162396. [PMID: 27647717](#).
- 7 Dhiwakar M, Clement WA, Supriya M, McKerrow WS. Antibiotics to reduce post-tonsillectomy morbidity. *Cochrane Database Syst Rev*. 2012;(2):CD005607. [PMID: 23235625](#).  
Mitchell RB, Archer SM, Ishman SL, Rosenfeld RM, Coles S, Finestone SA, et al. Clinical practice guideline: tonsillectomy in children (update). *Otolaryngol Head Neck Surg*. 2019;160(1 Suppl):S1–42. [PMID: 30921525](#).

- 8 Burton MJ, Glasziou PP, Chong LY, Venekamp RP. Tonsillectomy or adenotonsillectomy versus non-surgical treatment for chronic/recurrent acute tonsillitis. *Cochrane Database Syst Rev.* 2014;(11):CD001802. [PMID: 25407135](#).
- Francis DO, Chinnadurai S, Sathe NA, Morad A, Jordan AK, Krishnaswami S et al. Tonsillectomy for obstructive sleep-disordered breathing or recurrent throat infection in children. Rockville (MD): Agency for Healthcare Research and Quality (US); 2017;Report No.:16(17)-EHC042-EF. [PMID: 28182365](#).
- Mitchell RB, Archer SM, Ishman SL, Rosenfeld RM, Coles S, Finestone SA, et al. Clinical practice guideline: tonsillectomy in children (update). *Otolaryngol Head Neck Surg.* 2019;160(1):S1–42. [PMID: 30921525](#).
- 9 Brietzke SE, Shin JJ, Choi S, Lee JT, Parikh SR, Pena M et al. Clinical Consensus Statement: Pediatric Chronic Rhinosinusitis. *Otolaryngol Head Neck Surg.* 2014;151(4):542-53. [PMID: 25274375](#).
- Desrosiers M, Evans G, Keith P, Wright E, Kaplan A, Bouchard J, et al. Canadian clinical practice guidelines for acute and chronic rhinosinusitis. *Journal of Otolaryngology - Head & Neck Surgery.* 2011;40(2):S99-193. [PMID: 21310056](#).
- Shetty KR, Soh HH, Kahn C, Wang R, Shetty A, Brook C et al. Review and Analysis of Research Trends in Surgical Treatment of Pediatric Chronic Sinusitis. *Am J Rhinol Allergy.* 2020;34(3):428-35. [PMID: 31910642](#).
- Rosenfeld RM. Pilot Study of Outcomes in Pediatric Rhinosinusitis. *Arch Otolaryngol Head Neck Surg.* 1995;121(7):729-36. Web. [PMID: 7598848](#).

---

### **About the Canadian Society of Otolaryngology-Head & Neck Surgery**

The Canadian Society of Otolaryngology-Head & Neck Surgery (CSOHNS) is a proud partner of the Choosing Wisely Canada campaign. CSOHNS is an association that helps to serve the Canadian Otolaryngology-Head & Neck Surgery community. It is composed exclusively of otolaryngologists-head & neck surgeons and those training in the specialty. CSOHNS is dedicated to improving patient care through the support of education, the promotion of research, the dissemination of information, the scientific advancement of the Society, and the maintenance of high professional and ethical standards.



---

### **About Choosing Wisely Canada**

Choosing Wisely Canada is the national voice for reducing unnecessary tests and treatments in health care. One of its important functions is to help clinicians and patients engage in conversations that lead to smart and effective care choices.

🌐 [ChoosingWiselyCanada.org](http://ChoosingWiselyCanada.org) | ✉ [info@ChoosingWiselyCanada.org](mailto:info@ChoosingWiselyCanada.org) | 🐦 [@ChooseWiselyCA](https://twitter.com/ChooseWiselyCA) | 📘 [/ChoosingWiselyCanada](https://www.facebook.com/ChoosingWiselyCanada)