Personal Protective Measures in the Princess Margaret Head and Neck Cancer Clinic

Procedures will depend on the activity being performed and apply to symptom and travel free patients who have passed door screening:

- 1) Placing patients in the rooms and taking a history while sitting at least 3 M from the patient (as far as possible)
 - a. According to new UHN policy all health care workers working within clinical areas should wear a procedure mask (not N95) throughout their work period
- 2) Physical exam that involves trans-oral inspection and palpation
 - a. Gown, gloves, surgical mask and separate eye protection: goggles, visor or combination mask/visor
 - b. Donned and doffed in a sterile manner between each patient encounter
- 3) Nasopharyngoscopy
 - a. Full PPE gown, gloves, N95 mask and eye protection as above
 - b. Donned and doffed in sterile manner between each patient

PROCEDURES FOR FULL PPE DURING NASOPHARYNGOSCOPY

BEFORE PUTTING ON PPE:

- 1) Place garbage / linen bin in front of room
- 2) Place scope bin to send for re-processing outside of the room
- Remove your surgical mask (as per new UHN staff masking policy) and place in secure bag/locker etc.

ORDER OF DONNING PPE:

- 1) Tie hair back if it is long
- Thorough hand hygiene for at least 15 second using alcohol-based sanitizer preferably; if visibly soiled used soap and warm water
- 3) Gown first NOT the disposable yellow must be linen yellow put on the sleeves, tie the neck portion in front of you, pass over head, then tie behind
- 4) Then mask and eye protection (N95 with face shield surgical mask on top OR goggles OR visor)
- 5) Don gloves last. Gloves should cover the cuff of the gown.

CLOSE THE DOOR ONCE YOU ENTER THE ROOM

- 1) Perform the nasopharyngoscopy
- 2) Detach scope from light source and carry to door
- 3) Clinic Nurse stands outside door with gloved hands and holds scope bin
- 4) Place scope in bin without touching the sides or individual holding bin
- 5) Perform doffing procedure

ORDER OF DOFFING PPE: Dirtiest to cleanest

- 1) Gloves off first
- 2) Gown next untie from the back and take it off by rolling it into itself. Dispose of gown in the soiled linens bin.
- 3) Perform hand hygiene for minimum of 15 seconds with an Alcohol Based Hand Rub (soap and water if visibly soiled)
- 4) Remove eye protection (dispose of face shield if used, or remove goggles).
- 5) Remove Mask

- 6) Perform hand hygiene
- 7) Put on your assigned surgical mask from start of shift (unless visibly soiled or in direct contact with a positive/highly symptomatic patient)

Dr. John Waldron, Radiation Oncology